



BARENBRUG
agriseeds

PRODUCT GUIDE

The guide to Barenbrug Agriseeds
pasture cultivars and management.

Now is the perfect time to consider your pasture sowing options.

Good pasture is the cornerstone of New Zealand farming. You cannot buy a cheaper, more efficient and natural source of feed than your own grass and clover.

But not all pasture is equal. If you want to get the best out of your livestock and land, you need the right pasture to match your situation. Inside this guide you'll find the information you need to help select and grow superior pastures for your farm.



CONTENTS

Value of cultivar choice	4
Improving ryegrass persistence	5
Perennial ryegrass	6
Hybrid ryegrass	15
Italian and annual ryegrass	17
Brome grasses	20
Cocksfoot	22
White clover	24
Red clover	28
<i>AGRICOTE</i> seed treatment	30
Annual clovers (arrowleaf, balansa, Persian)	31
Chicory	34
Plantain	36
Brassica cultivars	38
Fodder beet	42
Ryegrass endophyte	44

VALUE OF CULTIVAR CHOICE

Introduction

Investing in the correct plant genetics is a key decision in pasture renewal. Cultivar choice affects a wide range of things including persistence, cool season yield, the heading date in grasses and late spring quality of a paddock and its palatability.

How do you value this?

One way is to look at an old ryegrass cultivar like *Nui*. At first glance it may look easier on the wallet than newer options but a quick cost:benefit analysis shows it is anything but.

It all comes down to performance. Comparative trials show a modern ryegrass like *Trojan* with *NEA2* endophyte will grow 2-3 t DM/ha a year more than *Nui*. On a sheep farm, that adds up to more ewes/ha, and more lambs too. On a dairy farm, *Trojan's* growth advantage means more milk in the vat. See the example below.

Bottom line? The correct cultivars, matched to the needs of the situation, pay for themselves surprisingly quickly, and after that, they're highly profitable.

New pasture genetics also have other benefits that old cultivars (like *Nui*) simply cannot match. Palatability and feed quality are higher, so animals perform better. Winter and early spring growth is much better, giving you more feed when you really need it. Endophytes enhance animal health and pasture persistence.

You wouldn't use a 30 year old ram or bull in your business because today's choices are so much more productive. Pasture is no different.

Example

Cost vs benefit: Modern ryegrass/white clover seed mix

Cost	
Typical modern seed	\$335/ha
Typical "cheap" seed	\$125/ha
Extra cost*	\$210/ha

* Estimated cost based on 20kg/ha ryegrass & 4kg/ha white clover.

Benefit - Sheep/beef	Benefit - Dairy
Extra 3 ewes/ha @ \$110 GM** = \$330	Extra 171kg MS/ha @ \$6 = \$1026/year ♦
Faster lamb growth (10% faster) from better pasture quality*** = \$62	Less cost production (30%***) = \$308/year
Extra benefit <u>each year</u> = \$392/ha	Extra benefit <u>each year</u> = \$718/ha

Assumptions:

**Extra 2 t DM/ha grown on sheep farm. Ewe gross margin (GM) = Income \$110/ewe (1.2 lambs @ \$90, cull ewes @ \$12, \$17.50 wool) less \$27.50/ewe costs.

*** Lamb LWG from Lincoln University trials at Ashley Dene, *Alto* ryegrass grew lambs 10% faster than *Nui* (over 5 separate 8 week periods). \$ benefit based on 15g/day faster LWG, 20 lambs/ha for 120 days = 36kgLW @\$1.70/kg = \$62.

♦ Extra milk solids (MS) based on 3 t DM/ha/year extra pasture; 80% utilisation; conversion to milk at 14kg DM/kgMS.

♦♦ Variable costs of extra MS produced vary farm to farm, but assumed 30% of income.




IMPROVING RYEGRASS PERSISTENCE

Introduction

The most commonly sown grass, ryegrass, needs to be well cared for to persist, and perform. If you look after it, it will look after you.

How to help pasture persist

1. Feed your pastures well. Soil phosphate and pH are particularly important for plant persistence. Ensure your soil fertility levels are correct, and keep them maintained.
2. Minimise plant death over summer. The most typical cause of plant death is a combination of moisture stress + overgrazing + insect feeding. For insect protection, use endophyte, combined with pasture cultivars bred for persistence. To avoid overgrazing in the dry a number of strategies help:
 - Sow summer crop (e.g. chicory, rape) to take pressure off pasture.
 - Have supplement on hand to feed out (or have access to supplement).
 - Plan ahead for key stock policy decisions to reduce feed demand (e.g. selling lambs store, milking once a day, selling trading stock).
 - Look after the best pastures - you can't look after every pasture, but look after your best and newest paddocks, as they'll grow fastest when rain comes.
3. Once it rains, don't change anything till pastures regrow. Grazing a pasture recovering from the dry too early can kill it – plants need 2-3 leaves per tiller so they have restored their reserves and will regrow quickly post-grazing.



4. Manage winter grazing to avoid damage from cattle. Spread stock out, or use a sacrifice area to keep them off your good paddocks. Use finer, denser ryegrass cultivars because they cover the ground and protect the soil better.
5. Consider other options. In very dry areas, alternative pastures may be better than ryegrass. Options include *Bareno* pasture brome and *Safin* cocksfoot.

TROJAN

PERENNIAL RYEGRASS

Trojan provides a balance of features previously unseen in a perennial ryegrass: exceptional DM yield across all seasons; excellent persistence; NEA2 endophyte for good animal health and insect control (including black beetle); high feed quality, and good resistance to rust and plant pulling.

High yield

Trojan NEA2 set a new standard of DM yield, but more important is the way it produces this yield at the shoulders of the season, in winter/early spring and summer.

Star FVI rating

Trojan NEA2 is in the top star category for DM yield for a diploid ryegrass in the DairyNZ Forage Value Index

In this example cultivars in this top group, which are ranked alphabetically, provide around \$413-\$527/ha more operating profit each year than the 1 Star cultivars like *Nui* ryegrass on a typical dairy farm.

DairyNZ FVI perennial ryegrass list 2019 - upper North Island*

FVI ¹ (Star rating)	FVI Star Rating (\$/ha)	Cultivar	Performance Values ² (1-5 rating)				
			Dry matter (DM)				
			Winter	Early spring	Late spring	Summer	Autumn
★★★★★	\$413 to \$527	Base AR37	5	4	4	5	5
		One50 AR37	5	4	4	5	5
		Platform AR37	5	5	4	5	5
		Prospect AR37	5	5	5	5	4
		SF Hustle AR1	5	5	5	5	4
		Trojan NEA2	5	5	5	5	4
★★★★★	\$298 to \$413	Excess AR37	5	5	4	5	5
		Governor AR37	4	4	4	5	5
		Halo AR37	5	4	3	4	4
		Matrix SE	4	5	4	5	5
		Raider NEA2	4	5	4	5	4
		Request AR37	5	5	5	4	4
		Ultra AR1	4	4	4	5	4
		★★★	\$184 to \$298	AberGreen AR1	3	4	5
Base AR1	4			4	3	3	3
Expo AR1	4			4	4	4	4
Expo AR37	5			4	4	4	4
Governor AR1	4			5	5	5	4
Ohau AR37	5			4	4	3	3
One50 AR1	4			4	3	4	4
Rely AR37	5			4	4	3	4
SF Moxie AR1	4			4	5	5	4
★★	\$69 to \$184	AberMagic AR1	4	2	4	4	4
		Excess AR1	3	4	4	4	4
		Rohan NEA2	4	3	2	3	4
		Samson SE	4	4	4	3	4
★	\$-45 to \$69	Bronte AR1	4	3	2	3	3
		Nui	3	4	3	2	3
		Pacific SE	3	4	4	2	3
		Samson AR37	4	4	3	3	4
	\$-375 to \$-45	AberGreen WE	1	1	1	2	2
		AberMagic WE	1	1	1	1	2
		Uncertif. P. Ryegrass	3	3	1	1	1

*The FVI lists are produced in February each year for four regions. Full lists at: www.dairynz.co.nz

Endophyte

Trojan contains *NEA2* endophyte, a mixture of different endophyte strains providing:

- Good control of adult black beetle (see page 44) and Argentine stem weevil.
- Very good control of pasture mealy bug (provisional rating).
- Moderate control of root aphid.

Feed quality

Trojan is late heading (+16 days) with a low level of aftermath heading, giving it better feed quality in late spring and summer. In trials for the DairyNZ Forage Value Index *Trojan* showed very high feed quality with an average 12.5 MJME/kg DM through the year.

Persistence

The persistence of *Trojan* has been excellent in trials. A couple of extreme tests of this were where *Trojan* persisted well in the Waikato, and recovered well after the severe 2007-08 summer drought. It also showed excellent persistence after five and a half years in the tough dryland Canterbury conditions at Mt. Possession, under average soil fertility and normal set stocking management with sheep.

Resistance

Trojan has good resistance to rust, and very good resistance to plant pulling.

Fine leaved

Trojan is medium-fine leaved and densely tillered.

Species classification

Trojan was bred as a perennial ryegrass and performs as one. It has a low level of tip awns (hairs) on its seed so under the seed certification regulations this means it is classified as *Lolium boucheanum*. In terms of pasture performance it is a perennial ryegrass.

Sowing *Trojan*

Dairy		kg/ha
Top performing palatable dairy pasture*	<i>Trojan</i> perennial ryegrass	18-22
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	Total	22-26
* <i>Captain</i> cool season plantain could be added to reduce N leaching & increase mineral content (see page 36)		
Sheep, Beef, Deer		kg/ha
Top performing, palatable pasture	<i>Trojan</i> perennial ryegrass	16-20
	<i>Weka</i> white clover	2
	<i>Apex</i> white clover	2
	<i>Safin</i> cocksfoot	2-3
	Total	22-27

ROHAN

SPREADING PERENNIAL RYEGRASS

Rohan spreading perennial ryegrass (*SPR*) was specifically bred to give sheep and beef farmers persistent, easy-care pasture. It has a unique spreading ability, very fine leaves and a dense habit. It has good palatability, with a tendency to stay leafy green and comes with *NEA2* endophyte for good animal performance.

Spreading habit

Rohan SPR has a spreading characteristic which provides two key advantages. First, it helps *Rohan SPR* fill bare areas in a pasture that may otherwise be occupied by weeds as shown in the photo below. This means *Rohan SPR* competes against weed ingress.

Second, *Rohan SPR's* spreading habit helps pastures recover from adverse climatic events, particularly extended dry periods, because it spreads to fill space where ryegrass tillers may have died.

Rohan SPR is not indestructible. Like all ryegrasses it will perform best under reasonable management and soil fertility. But its spreading habit improves persistence.



Rohan stolon spreading across the ground.

Sheep & beef system fit

The key with any pasture is matching a cultivar to a particular situation. So where does *Rohan SPR* fit? Looking at different pasture types across properties, based on persistence:

Pasture type	Finishing pasture	High yield, persistent pasture	Persistence key requirement	Toughest, non-ryegrass situations
	← <i>Increasing persistence</i> →			
Example	<i>Shogun Viscount</i>	<i>Trojan</i>	<i>Rohan SPR</i>	<i>Bareno Safin</i>
Description	High performance, palatable tetraploid ryegrasses are best suited for specialist finishing pastures.	<i>Trojan</i> provides an excellent balance of high DM yield and very good persistence that will suit many situations.	<i>Rohan SPR</i> takes things a step further, as a very persistent ryegrass suited to tougher conditions.	Some situations are just too tough or dry for ryegrass. This is where <i>Bareno</i> pasture brome and <i>Safin</i> cocksfoot suit.

**High yield
in dry
conditions**

Under ideal conditions *Rohan SPR* does not have the same yield potential as cultivars such as *Trojan*, producing about 10% less. But under tough, dry conditions *Rohan* will likely persist and yield more over the life of a pasture.

**'Easy care'
pasture**

Under semi-intensive to semi-extensive farm systems it is not easy to maintain pasture quality in late spring. Late heading cultivars help, but *Rohan SPR* is more than this, and in on-farm trials is showing 0.7 higher ME than some other cultivars in November and December. A continual comment from farmers with *Rohan SPR* is that it 'always looks good'. It stays greener and leafier and is often preferentially grazed.



Three year old Rohan SPR (green) sown beside Nui ryegrass (brown) in the same paddock in Central Otago. Rohan's been much better grazed by stock.

**NEA2 &
animal health**

Rohan SPR with *NEA2* endophyte provides very low staggers risk pasture for sheep and staggers free pasture for cattle. In the 15 years we have sold *NEA2*, no ryegrass staggers have been seen in sheep or cattle on commercial farms.

**Suggested
seed mix**

Sheep, Beef, Deer		kg/ha
Top performing, palatable pasture	<i>Rohan</i> perennial ryegrass	16-20
	<i>Weka</i> white clover	2
	<i>Apex</i> white clover	2
	<i>Safin</i> cocksfoot	2-3
Total		22-27

Sub clover(s) are often added to this mix.

Rohan Perennial Ryegrass is owned and marketed by Barenbrug Agriseeds
Rohan Perennial Ryegrass is protected under the NZ Plant Variety Rights Act 1987

TYSON

PERENNIAL RYEGRASS


Tyson is a leap forward in perennial ryegrass genetics for red meat farmers. It provides 35% more early spring feed, while maintaining high yield throughout the year, making it ideal for sheep and beef breeding operations.

Sheep breeding system fit

With its superb early spring growth *Tyson* can feed ewes better through early lactation, meaning better lamb growth, which in turn allows more lambs to be finished off mum. This has always been a key goal for sheep and beef farmers because:

- Early lambs meet early export schedules typically at better prices.
- Weaning check is avoided (2 weeks lost LWG) and dressing out percentage is higher.
- Extra feed is freed up for other stock.

Getting the best from *Tyson*



Cover target for lambing

While *Tyson* has the genetic potential to grow extremely well in early spring, it needs to be managed correctly to capture its full potential. It should be set stocked in spring at a **minimum** cover of 1200-1300 kg DM/ha (or 3-4 cm pasture height) for singles, 1500-1600 kg DM/ha (or 4-5 cm height) for twins, or 1700+ kg DM/ha for triplets through lambing.

Otherwise, *Tyson* won't have the leaves to capture enough sunlight to grow to its genetic potential, which is the science behind the saying 'grass grows grass.' Also as pasture height drops so does bite size, and although ewes take more bites both their pasture intake, and lamb growth rate, drop.

DM yield

Tyson has outstanding early spring growth in August and September, growing 35% more than the trial mean yield in inland Canterbury (altitude 190 m ASL). Total yield of *Tyson* over the year is excellent, similar to *Trojan*, with strong seasonal yield in summer and autumn.



The outstanding early growth of the three Tyson plots (marked by white lines in inset) stand out clearly in this trial. Photo taken in mid September in Canterbury 190m a.s.l.

Growth habit

Tyson is a fine leaved, densely tillered diploid perennial ryegrass suitable for both set stocking and rotational grazing.

Heading date

Tyson is the earliest heading of any perennial ryegrass on the market, with a -7 day heading date. For improved feed quality through the remainder of the season, *Tyson* was carefully selected for low aftermath heading.

Endophyte

Tyson is available with *ARI* endophyte, which provides very good control of Argentine stem weevil and pasture mealy bug, with no negative impacts on animal health. It's also available as a low endophyte option.

Suggested seed mix

Sheep, Beef, Deer		kg/ha
For systems requiring more early spring feed	<i>Tyson</i> perennial ryegrass	16-20
	<i>Apex</i> white clover	2
	<i>Weka</i> white clover	2
	<i>Morrow</i> red clover (coated)	6
	Total	26-30

Possible additions

- Timothy at 1 kg/ha – a highly digestible grass that suits summer moist areas.
- *Captain CSP* plantain at 2 kg/ha – provides extra summer feed value, lasts 2-3 years.

Tyson Perennial Ryegrass is owned and marketed by Barenbrug Agriseeds
Tyson Perennial Ryegrass is protected under the NZ Plant Variety Rights Act 1987


VISCOUNT

PERENNIAL RYEGRASS

Viscount is a standout tetraploid perennial ryegrass of the Barenbrug Agriseeds' plant breeding programme, with outstanding palatability, excellent early spring and total yield, improved rust resistance and high feed quality. *Viscount* has been upgraded to *NEA4* endophyte.

Seasonal growth

Viscount is late heading (+19 days). It has excellent early spring growth, coinciding with late calving or lambing when feed is most valuable. For dairy farmers extra grass at this time of year has been valued at an average of \$0.46/kg DM in the DairyNZ Forage Value Index.



Feed quality

Viscount provides high quality very palatable feed, with reduced aftermath heading and improved rust tolerance. It is more upright for ease of harvest, allowing good clover content and boosting animal performance.

NEA4 endophyte

Viscount comes with *NEA4* endophyte, which improves its persistence through better ASW, black beetle and root aphid control. *Viscount NEA4* provides excellent animal performance with a very low risk of animal health problems such as ryegrass staggers.

Suggested seed mixes

Dairy		kg/ha
For high feed quality and high yields	<i>Viscount</i> perennial ryegrass*	30
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	Total	34
Dairy, Sheep, Beef		kg/ha
For highly palatable pasture with extra robustness. (See page 13)	<i>Viscount</i> perennial ryegrass*	15
	<i>Trojan</i> perennial ryegrass	10
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	Total	29
Sheep, Beef		kg/ha
For high feed value pasture ideal for finishing	<i>Viscount</i> perennial ryegrass*	30
	<i>Weka</i> white clover	2
	<i>Apex</i> white clover	2
	<i>Morrow</i> red clover (coated)	6
	Total	40

* Tetraploids are sown at a higher rate than diploids, because of their larger seed.

MIXING TETRAPLOID & DIPLOID RYEGRASS


Summary

- Mixing *Viscount* with *Trojan* perennial ryegrass can produce higher animal performance with easier pasture management, than traditional pasture.
- This tetraploid/diploid mix fits a range of farm systems as it is much more persistent than a straight tetraploid pasture, because diploid plants protect the tetraploid.

Background

On many farms the tetraploid/diploid perennial ryegrass mix is now the norm, growing more energy (MJ ME/ha) and being easy to manage.

Tetraploid perennial ryegrasses, like *Viscount*, have excellent DM yield and year-round growth, but being so palatable, many farmers struggle to avoid over grazing them and getting the persistence they want. Adding a denser, finer diploid ryegrass to the mix changes the dynamics.




The tetraploid/diploid mix pasture is an average of the two types, denser and more robust than a straight tetraploid, and much more palatable than a straight diploid pasture.

Palatability & stems

The palatability of the pasture is due to *Viscount's* soft stems, which give improved animal performance and hold quality even at high covers (e.g. 3500-3600 kg DM/ha) making them very easy to graze.

While straight tetraploid pastures are often overgrazed, reducing persistence, in the mix they are protected by the denser, finer, diploid plants.

Diploid protects from overgrazing



Pre-grazing

Tetraploid plants (dark green) & diploid (light green) are mixed up.



Post-grazing

Tougher diploid stems help protect tetraploid plants from over-grazing.

Sowing rate

We have tested different tetraploid/diploid perennial ryegrass mixes and recommend sowing half the normal rate of each cultivar, e.g. 15 kg/ha of tetraploid *Viscount* (half of 30 kg) plus 10 kg/ha of diploid *Trojan* (half 20 kg).

GOVERNOR

PERENNIAL RYEGRASS

Governor combines genetics from two of Barenbrug Agriseeds' most popular previous cultivars to set a new standard for an all-round pasture. With outstanding survival and excellent DM yield on the shoulders of the season.

Genetic legacy

The persistence of *Bronsyn*, with the high DM yield and palatability of *Tolosa*, make *Governor* ideal for dairy, sheep and beef systems.

Persistent

Governor has shown outstanding survival through drought and high insect pressure under grazing in farm trials across the country. Fine, densely tillered and diploid, we believe it is the premium AR37 cultivar of choice for persistence. It is also available with ARI endophyte for parts of the lower North Island and the South Island where AR37 isn't required.

Seasonal growth

A key feature is *Governor's* ability to grow more DM on the shoulders of the season, in early spring and autumn, when it is most needed.

All-rounder

With a +5 days heading date, low aftermath heading (similar to *Alto*) and better rust resistance than its parents, *Governor* is the reliable, persistent all-rounder.

Where to sow

Pasture type	High animal performance	High yield, persistent pasture	More density & robustness	Persistence key requirement	Toughest, non-ryegrass situations
Example	<i>Trojan/Viscount Mix</i>	<i>Trojan</i>	<i>Governor</i> OR <i>Trojan/Rohan</i>	<i>Rohan SPR</i>	<i>Bareno Safin</i>
Description	<i>Trojan</i> provides density and robustness, tetraploid <i>Viscount</i> adds high palatability driving animal intakes.	<i>Trojan</i> provides an excellent balance of high DM yield and very good persistence, that will suit many situations.	<i>Governor</i> is a fine, dense cultivar and provides a robust pasture with AR37 or ARI endophyte. Mixing <i>Rohan</i> with <i>Trojan</i> gives a similar dense, robust pasture.	<i>Rohan SPR</i> is a very persistent spreading ryegrass suited to hill country and tougher conditions.	Some situations are just too tough or dry for ryegrass. This is where <i>Bareno</i> pasture brome and <i>Safin</i> cocksfoot suit.

Suggested seed mix


Dairy	kg/ha
Top performing palatable dairy pasture	<i>Governor</i> perennial ryegrass 18-22 <i>Kotuku</i> white clover 2 <i>Weka</i> white clover 2 <hr/> Total 22-26
Sheep, Beef, Deer	kg/ha
Top performing, palatable pasture	<i>Governor</i> perennial ryegrass 16-20 <i>Weka</i> white clover 2 <i>Apex</i> white clover 2 <i>Safin</i> cocksfoot 2-3 <hr/> Total 22-27

SHOGUN

HYBRID RYEGRASS

Shogun created a new position in the market for hybrid ryegrass. Winter growth is equal to many Italian ryegrasses, and *Shogun* outyields many perennials during summer. Persistence is excellent for a hybrid, with *NEA* endophyte for insect protection.

Shogun redefines ryegrass categories



High yield

In trials *Shogun* has significantly out-yielded other hybrid cultivars. As well as excellent cool season growth, it has exceptional summer and autumn yield.

Hybrid Ryegrass: All New Zealand trials (yields by season as percentage of Mean and LSI)

Entry	Number of Trials	Winter		Early Spring		Late Spring		Summer		Autumn		Total	
		% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI
<i>Shogun NEA</i>	11	117.1	6.3	107.7	4.4	106.6	3.4	113.4	4.0	108.4	6.6	109.7	3.8
<i>Ohau AR37</i>	5	93.4	9.9	100.3	6.8	99.9	5.3	95.3	6.2	107.0	10.3	99.6	5.9
<i>Ohau AR1</i>	7	88.5	8.4	98.5	5.8	102.0	4.5	95.4	5.3	93.4	8.7	97.0	5.0
<i>Asset AR37</i>	8	100.9	7.9	93.6	5.4	91.5	4.2	96.0	4.9	91.2	8.2	93.7	4.7
Mean (kg DM/ha)	12	898		1878		3109		2571		1875		10332	

NFVT Summary 1991 – 2018 (August 2018). If two means differ by more than the sum of their least significant intervals (LSI), they are significantly different at the 5% level

Fast establishment

Shogun's fast establishment is comparable to that of an Italian ryegrass. This allows paddocks resown with *Shogun* to be brought back into the grazing rotation more quickly than those renewed with perennial or other hybrid ryegrasses.

Black beetle control

Shogun with *NEA* endophyte has good control of black beetle, equal to *Viscount NEA4* or *Trojan NEA2*.

2-5 year option

Under good grazing management *Shogun* is a 2-3 year option in summer dry areas, 3-5 years in summer moist. Persistence is aided by its *NEA* endophyte.

Great animal health

NEA endophyte is one of the most animal safe endophytes available. However, there is a low risk of *NEA* endophyte causing a low level of ryegrass staggers in sheep or deer in extreme situations (where animals are forced to graze right into the base of a pasture in very summer dry conditions).

Feed quality

Shogun has excellent summer quality, with a very late heading date (+26 days) and little aftermath heading or seeding through the summer.

Palatability

When sheep or cattle like a pasture, they eat more of it, and liveweight gains (LWG) increase. *Shogun* delivers high animal intakes and growth rates.

Suggested seed mixes

Dairy		kg/ha
High performance 3-5 year pasture	<i>Shogun</i> hybrid ryegrass	30
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	Total	34
Undersowing as fast establishing 2-3 year pasture (with black beetle control)	<i>Shogun</i> hybrid ryegrass	13-20*
	Total	13-20*
Sheep, Beef, Deer		kg/ha
High performance finishing pasture	<i>Shogun</i> hybrid ryegrass	30
	<i>Weka</i> white clover	2
	<i>Apex</i> white clover	2
	<i>Morrow</i> red clover (coated)	6
	Total	40

*Sowing rate depends on how thin pasture to be undersown is. Tetraploids are sown at a higher rate than diploids, because of their larger seed.



*Extreme palatability differences in animal health trials in spring, when there were no fences between plots set stocked with lambing ewes. *Shogun* NEA is front left and back right; SE perennial ryegrass is front right and back left.*

TABU+

ITALIAN RYEGRASS

Tabu+ was bred to supersede the best selling *Tabu*, with significantly higher total DM yield. *Tabu+* is nutritious, with explosive establishment speed and superior cool season growth.

Multi-use

Tabu+ is suitable as an 8-12 month high performance crop; can last 2-3 years in areas with mild summers, or can be used for undersowing into run out pasture to boost winter-spring growth. In dense pastures spraying before drilling is recommended.

Note: In situations where a pasture is required for 18+ months *Shogun NEA* may be a better option.

High DM yield

Tabu+ is the top yielding Italian ryegrass in the National Forage Variety Trials (NFVT) with significantly more winter growth. It is out yielded only by *Shogun NEA* hybrid ryegrass.

Italian Ryegrass: All New Zealand trials (yields by season as a percentage of Mean and LSI)

Entry	Number of Trials	Establishment Autumn		Winter		Early Spring		Late Spring		Summer		Total	
		% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI
<i>Shogun NEA</i>	7	97.2	6.3	96.5	5.3	106.6	4.4	113.2	4.6	124.0	7.9	110.8	3.8
<i>Tabu+ WE</i>	6	105.7	6.7	110.5	5.7	105.4	4.7	106.0	4.9	111.4	8.4	107.9	4.1
<i>Asset AR37</i>	18	100.6	4.0	103.1	3.4	98.0	2.8	98.3	2.9	111.9	5.0	102.8	2.5
<i>Supercruise WE</i>	10	106.8	5.5	106.5	4.6	96.8	3.8	103.3	4.0	102.7	6.8	102.6	3.3
<i>Lush AR37</i>	12	106.3	4.9	102.4	4.1	99.0	3.4	94.5	3.6	106.4	6.1	101.0	3.0
<i>Jackpot WE</i>	7	99.7	6.3	101.4	5.3	100.5	4.4	100.8	4.6	99.9	7.9	100.4	3.8
<i>Vibe WE</i>	6	106.0	6.8	99.0	5.8	95.3	4.8	101.5	5.0	99.6	8.5	100.0	4.2
<i>Feast II WE</i>	36	98.8	2.8	98.8	2.4	99.3	2.0	98.7	2.1	98.8	3.5	98.9	1.7
<i>Blade WE</i>	10	104.9	5.4	99.8	4.5	101.6	3.8	97.3	3.9	93.1	6.7	98.2	3.3
<i>Mona WE</i>	9	97.8	5.7	97.6	4.8	102.2	4.0	100.0	4.1	93.0	7.1	98.0	3.4
<i>Asset WE</i>	6	94.5	6.7	96.7	5.6	96.5	4.7	98.9	4.8	99.7	8.3	97.8	4.1
<i>Sonik WE</i>	9	96.3	5.6	99.3	4.7	102.1	3.9	97.9	4.0	92.8	6.9	97.3	3.4
<i>Moata WE</i>	24	85.3	3.5	88.4	2.9	96.7	2.5	89.6	2.5	66.6	4.4	84.2	2.1
Mean (kg DM/ha)	77	1710		1787		2948		4135		3894		14474	

NFVT Summary 1991 – 2018 (August 2018)

LSI (Least Significant Interval) - If two means differ by more than the sum of their LSI, they are significantly different at the 5% level

2000% ROI

Tabu+ produced an extra 340 kg DM/ha over *Tabu* as an 8 month crop (from sowing through winter to mid spring). For a small additional seed cost (e.g. \$5/ha) for *Tabu+*, that equates to a 2000% ROI, given this cool season feed is highly valuable, i.e. \$0.35/kg DM or an extra \$103/ha (value calculated from the DairyNZ FVI).

Sowing *Tabu+*

Winter ryegrass crop	kg/ha
<i>Tabu+</i> Italian ryegrass	18-22
Winter ryegrass crop with annual clovers	kg/ha
<i>Tabu+</i> Italian ryegrass	18-22
<i>Laser</i> Persian clover	4
<i>Vista</i> balansa clover	3
Total	25-29
2-3 year pasture option	kg/ha
<i>Tabu+</i> Italian ryegrass	18-22
<i>Morrow</i> red clover (coated)	6
<i>Kotuku</i> or <i>Apex</i> white clover	2
<i>Weka</i> white clover	2
Total	28-32
Undersowing	kg/ha
<i>Tabu+</i> Italian ryegrass	10-15*
<i>Kotuku</i> or <i>Apex</i> white clover	1.5
<i>Weka</i> white clover	1.5
Total	13-18

*Sowing rate varies depending on how thin pasture to be undersown is.



Tabu+ has explosive establishment speed and cool season growth.

HOGAN

ANNUAL RYEGRASS

Hogan sets a new standard for annual ryegrass. It produces 1 t DM/ha more than old, widely sown cultivar *Tama*, extra growth which is worth up to \$380/ha.

High value

Hogan establishes rapidly and out produces 30+ year old *Tama* by 1 t DM/ha. *Hogan's* advantage is valued by the 2019 DairyNZ Forage Value Index at \$380/ha extra profit, i.e. a 10 fold return on the extra \$35-\$45/ha it costs to sow *Hogan* over *Tama*.

Fast establishment

Hogan is a tetraploid bred for rapid establishment (21% faster than *Tama*) to provide fast feed in autumn, a critical advantage particularly following dry summers.


Hogan is in the top ranking for annual ryegrass in the National Forage Variety Trials.

Annual Ryegrass: All New Zealand trials (yields by season as percentage of Mean and LSI)

Entry	Number of Trials	Establishment Autumn		Winter		Early Spring		Late Spring		Total	
		% of Mean	LSI	% of Mean	LSI	% of Mean	LSI	% of Mean	LSI		
<i>Hogan WE</i>	7	108.4	6.6	106.1	5.4	100.3	4.5	106.9	5.2	105.1	3.6
<i>Dash WE</i>	6	99.0	7.0	100.5	5.8	108.4	4.8	107.2	5.6	105.0	3.9
<i>Zoom WE</i>	5	99.9	7.8	102.4	6.4	100.7	5.3	105.0	6.1	102.5	4.3
<i>Winter Star II WE</i>	7	102.8	6.5	102.8	5.3	102.6	4.4	101.6	5.1	102.3	3.6
<i>Tama WE</i>	17	87.2	4.1	97.3	3.4	94.7	2.8	92.3	3.2	93.0	2.3
<i>Progrow WE</i>	9	102.6	5.9	90.9	4.9	93.3	4.1	87.0	4.7	92.1	3.3
Mean (kg DM/ha)	77	1654		1765		2923		3677		10019	

NFVT Summary 1991 – 2018 (August 2018)

If two means differ by more than the sum of their least significant intervals (LSI), they are significantly different at the 5% level



Suggested seed mixes

Hogan annual ryegrass can be sown alone, or mixed with oats or turnips.

Dairy, Sheep, Beef, Deer	kg/ha
For winter crop	<i>Hogan</i> annual ryegrass* 30
	Total 30
Winter ryegrass crop with annual clovers	<i>Hogan</i> annual ryegrass 26-30
	<i>Laser</i> Persian clover 4
	<i>Vista</i> balansa clover 3
	Total 33-37

* Tetraploids are sown at a higher rate than diploids, because of their larger seed.

Hogan annual ryegrass is owned and marketed by Barenbrug Agriseeds
Hogan annual ryegrass is protected under the NZ Plant Variety Rights Act 1987

BARENO

BROME

Bareno is persistent, high yielding and very palatable. Paddocks of *Bareno* are assets in a dryland farm system as they persist much better than perennial ryegrass, tolerating dry conditions and hard grazing.

Easy management

Bareno is quite different to other bromes, and is more flexible in its management. It is more persistent than prairie grass, and can tolerate both rotational grazing and set stocking. It should be noted that the persistence of *Bareno* may decline north of Taupo, therefore it is not as suited to northern North Island areas.

Highly palatable


A strong feature of *Bareno* is its palatability through all seasons:

- It remains palatable, even when seed heads are present, and stays greener and leafier than ryegrass in summer.
- *Bareno* can support a high legume content (40% sub and white clover has been measured in spring).

High yield

On Lincoln University's Silverwood Farm, a dryland breeding property in inland Canterbury, *Bareno* produced 12.5 t DM/ha/year, 9% more than new sowings of perennial ryegrass (11.5 t DM/ha), with excellent spring, summer and autumn growth.

Seasonal growth



Sow early

Brome grasses are slower to establish than ryegrass, so make sure to:

- Sow when warm - soil temperature 12°C+.
- Prepare a good seedbed, preferably using a summer fallow.

Suggested seed mixes

Sheep, Beef, Deer	kg/ha
Persistent dryland pasture	<i>Bareno</i> pasture brome 25-32*
Can be added:	
<i>Safin</i> cocksfoot	Inclusion of species depends on situation. Seek advice if unsure.
Sub clover	
<i>Apex</i> white clover	
<i>Morrow</i> red clover	

**Bareno* sowing rate high because brome grasses have large seeds.

BARENO MANAGEMENT

Growing *Bareno*

Brome grasses are slower to establish than ryegrass. If you spend a little extra time on correct sowing and early management, you'll be rewarded with good results.

Preparation

A summer fallow prior to late summer/early autumn sowing is the recommended best practice to establish *Bareno* pasture in dry areas. This allows moisture to be carried from the spring through to sowing, ensuring good results even in a dry autumn. Prepare a run-out paddock by spraying out or cultivating in spring (Oct/Nov) before pastures dry out. If there is a further weed strike, spray again before drilling.

If cultivating, prepare a fine, well compacted, weed-free seed bed to allow correct seed depth and soil moisture retention for fast germination. Direct drilling has proven to be very successful. This fits well with summer fallow management.

Timing

Bareno is best sown when soil temperatures are above 12°C, during late summer or early autumn. This gives plants time to adequately establish before winter. Establishment is much slower in cool conditions.

Drilling

Sow seed shallow, at 10-20 mm. Take care when drilling - the seed may not flow well through some drills.

Managing *Bareno*

Bareno should not be overgrazed in its first year to allow plants to fully establish. *Bareno* can set seed quickly, however seed heads are much more palatable than those of other pasture grasses.

In dry summer conditions, *Bareno* pastures should not be bared out (although they will tolerate this better than ryegrass). Post-grazing covers of 3-4 cm will ensure persistence and regrowth through summer. Remember the plant's reserves in grasses are above the ground (not in the roots).



Bareno pasture brome is a persistent, easy to manage pasture, seen here at Mt. Possession in Canterbury.

SAFIN

SUPERFINE COCKSFOOT

Safin is an innovative super-fine leaved cocksfoot which will change farmers' perceptions about this type of grass. Traditional cocksfoot gained a bad reputation for becoming clumpy and unpalatable, eventually dominating swards. *Safin* sets a new standard, looking almost as fine as ryegrass. This is an exciting pasture development for dryland farmers in particular.

Early growth (with high total DM)

A key feature of *Safin* is its increased production in winter and early spring. DM growth is critical through lambing or calving for dryland farming systems, to finish stock prior to potential summer dry conditions. *Safin* has an advantage through this period, as shown below, and in the paddock it is noticeably faster to get away in spring.

Over the whole year total DM production of *Safin* is very good.

Cocksfoot yields in Canterbury*

Entry	Winter		Early spring		Late spring	Summer	Autumn	Total
<i>Safin</i>	123	a	124	a	104	105	119	110
<i>Ella</i>	90	b	101	b	100	114	111	106
<i>Wana</i>	82	b	117	ab	96	106	113	104
<i>Vision</i>	96	ab	108	ab	106	98	95	102
<i>Kara</i>	109	ab	107	ab	95	105	98	102
LSD (5%)	32		22		10	17	41	13

* Combined analysis of 2 trials run on Barenbrug Agriseeds Research farm, Courtenay, between 2004 and 2009.

Tiller density

Safin is a 'superfine' cocksfoot, which in fact looks very similar to ryegrass. It is finer and denser than other cocksfoots on the market, with significantly more tillers as shown in the table below. On its release *Ella* was considered fine leaved, but *Safin* sets a new benchmark, with 41% more tillers than *Ella*.

Cocksfoot tiller density in Canterbury dryland grazing trial

Entry	Tillers/m ²	Relative to <i>Ella</i> at 100%
<i>Safin</i>	431 a	141 %
<i>Ella</i>	305 b	100 %
<i>Tekapo</i>	303 b	91 %
<i>Greenly</i>	270 b	81 %
<i>Vision</i>	270 b	81 %
<i>Kara</i>	245 b	73 %
Trial mean	334	95
LSD (5%)	70	23

Pest control

Once established *Safin* is tolerant to grass grub and Argentine stem weevil (ASW) attack. However, seedlings are susceptible at sowing and *AGRICOTE Grass* seed treatment is recommended where these pests are a risk.

Cocksfoot has no endophyte and is therefore safe to graze low over summer.

Managing *Safin*

Cocksfoot is slower to establish than ryegrass. In a ryegrass-based pasture little cocksfoot is often seen until the first summer after it has been sown, after which the content of cocksfoot will increase, especially under dry conditions.

If sowing a specialist cocksfoot-based pasture, sow early while soil temperatures are above 12°C. A summer fallow, to conserve moisture prior to a late summer/early autumn sowing, is recommended in (potentially) autumn dry conditions.

Cocksfoot is lower in feed value and palatability than ryegrass if it becomes long or rank, so it should be kept short and leafy through spring. Graze cocksfoot when it has 3-4 leaves/tiller. The feed value of cocksfoot declines if it grows to 5 leaves/tiller, as older leaves die. In grazing trials, animal performance is reasonably good where cocksfoot has been kept leafy. Maintaining good legume content in cocksfoot pastures will also improve animal performance.

Cocksfoot is more tolerant of low-moderate soil fertility than perennial ryegrass. However, it is very responsive to nitrogen fertiliser, generally at higher response rates than ryegrass, which can be a very useful tool to promote growth, as well as improving feed value and protein content.

Suggested seed mixes

Sheep, Beef, Deer		kg/ha
As component of pasture mix	<i>Safin</i> cocksfoot	3-6
For cocksfoot-based pasture	<i>Safin</i> Cocksfoot	8-10
	Sub clover	6-8
	<i>Apex</i> white clover	2
	<i>Weka</i> white clover	2
	Total	18-22



Older, traditional cocksfoot (left) can form unpalatable clumps in a pasture, compared to superfine *Safin* cocksfoot (right).

KOTUKU

WHITE CLOVER

Kotuku is a very high yielding large leaved white clover with superior summer growth. It establishes well, has good persistence, and suits both dairying and sheep/beef finishing systems.

Why *Kotuku*?

White clover is critical for nutritive value and N fixation in pastures. It is also an important source of protein and ME for milking and growing stock, particularly in summer. *Kotuku* shows excellent seasonal growth, and outperforms all other trialed cultivars over the critical summer period.

High yield

This mixed sward trial included one entry without clover (no clover). The effect of clover on N fixation and yield is seen in the trial, with *Kotuku* showing particularly good yield due to its compatibility with ryegrass. Every extra 1t DM/ha of clover provides 25-30 kg N/ha 'free' nitrogen.

Seasonal DM yield data 2013-2016, Courtenay, Canterbury. Trial mean = 100.

Entry	Autumn	Winter	Early Spring	Late Spring	Summer	Total
<i>Kotuku</i>	117 a	107 bc	107 ab	109 a	121 a	114 a
<i>Kopu II</i>	114 a	115 a	112 a	109 a	110 b	111 ab
<i>Kotare</i>	105 bc	106 c	108 a	111 a	108 bc	106 bc
<i>Tribute</i>	102 bc	105 c	109 a	107 ab	102 bd	105 c
<i>SF Quest</i>	106 b	114 ab	111 a	105 ac	98 d	104 cd
<i>Mainstay</i>	110 ab	101 cd	100 c	99 bd	102 bd	102 cd
<i>Weka</i>	99 cd	97 de	100 bc	106 ac	101 cd	100 de
<i>Bounty</i>	94 de	88 f	92 d	97 cd	102 cd	97 e
<i>Huia</i>	88 e	91 ef	92 d	95 d	97 d	95 e
No clover	46 f	67 g	59 e	55 e	29 e	44 f
Trial mean (kgDM/ha)	1765	721	970	1659	3101	8509
Significance	***	***	***	***	***	***

*Data from Courtney, Canterbury, 2013-2016. Statistical significance lettering is given, yields with the same letter are not significantly different at the 5% LSD level.



Persistence

Kotuku has demonstrated robust persistence for a large leaved cultivar, and has a medium stolon density. This makes it well-suited to driving summer production in dairying and dry stock finishing systems.

Fast establishment

Kotuku has consistently shown fast establishment. This can assist with broadleaf herbicide applications, where new clovers need to be at the 3-4 trifoliolate leaf stage.



Establishment speed of Kotuku (left) versus Mainstay.

Suggested seed mixes

Dairy		kg/ha
Top performing palatable dairy pasture	<i>Trojan NEA2</i> perennial ryegrass	18-22
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	Total	22-26
Sheep, Beef & Deer		kg/ha
High feed tetraploid pasture for finishing	<i>Viscount NEA4</i> perennial ryegrass	30
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	<i>Morrow</i> red clover (coated)	6
	Total	40

WEKA

WHITE CLOVER

Weka is a medium leaved high yielding white clover suited to all grazing systems. It has a high stolon density, a strong spreading habit, and excellent tolerance to clover root weevil (CRW).

High yield

In trials *Weka* has shown very high total yield, with good growth in all seasons, particularly through autumn and winter. Every extra 1t DM/ha of clover provides 25-30 kg N/ha 'free' nitrogen.

Yield scores of medium-large leaved clovers 2003-08*

Cultivar	Autumn	Winter	Early spring	Late spring	Summer	Total
<i>Weka</i>	6.1 a	5.2 a	5.8 a	6.0 a	5.8 a	5.8 a
<i>Tribute</i>	5.6 ab	4.4 a	5.7 a	5.9 a	5.8 a	5.5 a
<i>Sustain</i>	4.6 b	4.5 a	4.8 b	5.2 b	5.2 a	5.0 b

*Combines yield scores over 4 trials in the Waikato 2004-08 & 2005-08, & Canterbury 2003-06 & 2005-08. Yield scored on 1-9 basis, where 9 = very high yield. Yields with same letter not significantly different at 5% LSD level.

Persistence

Weka has excellent persistence because of its high tolerance to CRW, dry conditions, pugging and hard grazing, and its strong spreading habit. Yield scores under high levels of CRW attack show *Weka* is a very good choice for these conditions.

Yield of medium-large leaved clovers under CRW attack*

Cultivar	Yield under CRW attack
<i>Weka</i>	5.5 a
<i>Tribute</i>	4.9 b
<i>Sustain</i>	4.1 c

*Combines 5 yield scores over 2 Waikato trials in 2006 & 2007 in periods of high CRW damage. Yield scored 1-9, where 9 = very high yield. Yields with same letter not significantly different at 5% LSD level.

Suggested seed mix

All systems	kg/ha	
Productive, persistent clover combination	Perennial ryegrass	18-30
	<i>Weka</i> white clover	2
	<i>Kotuku</i> or <i>Apex</i> white clover	2
	Total	22-34



Weka's strong spreading habit and high CRW tolerance help it persist.

APEX

WHITE CLOVER

Apex is a breakthrough in persistent high yielding clover, adapted to summer dry conditions, with good clover root weevil tolerance.

Medium small leaf size

Apex has a medium-small leaf size, with significantly more stolon growing points than traditional cultivars like *Huia*, for improved drought and pest tolerance.

Good persistence

A key feature of *Apex* is its improved persistence. A four year trial under grazing in the Waikato showed *Apex* has excellent persistence into the fourth year, with the highest fourth year yields.

High yield

Apex has shown high yields in sheep grazing trials, particularly in the winter-spring and autumn periods.

Yield in three Manawatu sheep grazing trials (*Huia* = 100)*

Cultivar	Winter	Spring	Summer	Autumn
<i>Apex</i>	156 a	139 a	108 a	127 a
<i>Huia</i>	100 b	100 b	100 a	100 b
LSD (5%)	22	18	15	18

* Woodfield et al. NZ Grassland Association 63: 103-108

Spreading growth



Apex spreads strongly across bare ground, increasing legume content.

Suggested seed mix

Sheep, Beef & Deer		kg/ha
For more clover in grazing systems	Perennial ryegrass	18-30
	<i>Apex</i> white clover	2
	<i>Weka</i> white clover	2
	Total	22-34

Apex White Clover is marketed by Barenbrug Agriseeds
Apex was bred under a cooperative programme with AgResearch Grasslands
Apex White Clover is protected under the NZ Plant Variety Rights Act 1987

MORROW

RED CLOVER

Morrow comes from a tough family. Most red clovers wouldn't last long under intensive rotational dairy grazing on light, summer dry upper North Island soils. But *Morrow's* parents did. Over time, they adapted, survived, and stood out as good growers, even after repeated droughts.

We took plants from these old pastures, and selected the best of them for high yield, persistence and flowering to create a game-changing, multi-stemmed red clover for improved production and persistence under grazing.

Longevity

Red clover's biggest drawback has always been limited persistence under grazing. *Morrow's* improved grazing tolerance – helped by its semi-prostrate form and high stem count – means it will keep boosting production year on year. Like all red clovers *Morrow* will persist best on free-draining soils under a longer summer grazing round.

Yield + quality when it counts

High ME and high DM together create ideal late spring and summer finishing feed, giving high quality as grass ME drops off, and driving rapid liveweight gains for lambs and cattle.

Red clover seasonal yield in dryland Canterbury*

Entry	Early Spring	Late Spring	Summer	Autumn	Winter	Total yield
<i>Morrow</i>	6.4 a	7.0 a	7.4 a	7.7 a	5.4 a	6.8 a
<i>Tuscan</i>	6.1 ab	6.7 a	5.5 b	6.0 ab	6.1 a	6.1 ab
<i>Rossi</i>	5.2 ac	5.7 ab	5.3 bc	6.3 ab	5.5 a	5.5 b
<i>Relish</i>	4.3 c	4.3 b	3.7 c	5.3 b	3.0 b	4.0 c
Trial mean (kg DM/ha)	4.9	5.6	5.4	5.8	4.4	5.3
%CV	20.1	18.0	16.5	18.4	21.3	12.8

*Data from 2 years of pure sward trial, grazed by sheep. Trial sown 2016. Yield visually scored on 1-9 basis, where 9 = highest yield.



Morrow's excellent summer yield and feed quality make it ideal for stock finishing

CRW tolerance

Clover root weevil remains a pest of white clover throughout NZ, particularly in Northland where the biocontrol wasp hasn't established. Red clover is tolerant of clover root weevil, providing pasture species diversity and extra legume content.

Free N

Red clover fixes its own nitrogen, adding it naturally to pastures in a slow, continual way. This will be ever more important to provide N on farm, as fertiliser comes under increasing environmental scrutiny. *Morrow* can fix over 200 kgN/ha based on its yield (about 25 kgN per t DM grown).

Phyto-oestrogen levels

Morrow has medium oestrogen levels. This means care needs to be taken to avoid grazing when mating ewes or hoggets, 3-6 weeks either side of mating.

Suggested seed mix

All systems		kg/ha
Productive, persistent clover combination	Perennial ryegrass	18-30
	<i>Morrow</i> red clover (coated)	6
	<i>Kotuku</i> or <i>Apex</i> white clover	2
	Total	26-38
Sheep, Beef, Deer		kg/ha
Two year high LWG finishing crop	<i>Captain</i> plantain	10
	<i>Laser</i> Persian clover	4
	<i>Vista</i> balansa clover	3
	<i>Morrow</i> red clover (coated)	6
	<i>Weka</i> white clover	4
Total	27	
Perennial ryegrass - Finishing		kg/ha
Fanatastic feed quality combined with animal performance.	<i>Viscount</i> tetraploid perennial ryegrass	16
	<i>Shogun</i> tetraploid perennial ryegrass	10
	<i>Weka</i> white clover	2
	<i>Apex</i> white clover	2
	<i>Morrow</i> red clover (coated)	6
	<i>Captain</i> plantain	2
Total	38	

AGRICOTE SEED TREATMENT

AGRICOTE helps ensure good even establishment of new pastures and crops.

Best possible start

Establishment is a critical time for a new pasture or crop as its potential performance is determined in this early stage. Different *AGRICOTE* seed coatings aid establishment by helping protect your seedling plants from insects and fungal diseases and by supplying nutrients to clovers.

If you have a pasture or crop that fails, the main cost (usually 75-80%) is the lost feed. There is also an additional cost in resowing as the example below shows.

Example - cost of a pasture failure.

What happens	Autumn pasture fails to establish
Cost of lost DM	Loss of 5 t DM/ha production (from April – September) = \$1500/ha (valued at 30c/kg DM*)
Cost of resowing spring	= \$500/ha (to re-spray, light cultivation, buy seed & resow)
Total cost failure	= \$2000/ha

*April – September pasture has a high value of 30c/kg DM as this is a feed deficit period.

Order your seed with *AGRICOTE* coating to get the following (Y=Yes):

Grass seed treatment

Seed treatment	Insect protection			Fungal pathogens		Other	Sowing rate
	Argentine stem weevil	Black beetle	Grass grub	Fusarium	Pythium	Weight build up	
<i>AGRICOTE GRASS</i>	Y	Y	Y	Y	Y	Nil	Same as bare

Clover seed treatment

Seed treatment	Insect protection	Fungal pathogens			Additives		Other	Sowing rate
	Nematodes	Fusarium	Pythium	Rhizoctonia	Lime	Nutrients	Weight build up	
<i>AGRICOTE CLOVER</i>	Y	Y	Y	Y	Y	N, P, Mn, Zn, Mo	75%	4kg*

Brassica seed treatment

Seed treatment	Insect protection			Fungal pathogens		Additives	Other	Sowing rate
	Nysius	Spring tails	Aphids	Fusarium	Pythium	Molybdenum	Weight build up	
<i>AGRICOTE BRASSICA</i>	Y	Y	Y	Y	Y	Y	Nil	Same as bare

ZULU II

ARROWLEAF CLOVER

Zulu II is mid to late flowering and produces high ME feed for grazing, finishing or silage from early spring through to early summer. It has a deep tap-root to aid summer growth; hard seed and good regeneration if managed well. *Zulu II* is also showing very good potential as a productive alternative to sub clover in hill country.

Yield + quality

Zulu II can transform low-yielding dry paddocks into palatable, productive pastures growing over 10 t DM/ha in spring and early summer. Feed value is excellent, with less risk of bloat than other annual clovers.

System fit

For dairying – particularly on free-draining dryland – add *Zulu II* to Italian ryegrass to extend late spring and summer growth and improve pasture ME. For sheep and beef, sow a straight sward; mix with plantain and other clovers or oversow into existing hill pastures. *Zulu II* best suits rotational grazing.

Management

For persistence in hill country *Zulu II* is hard seeded and will not need to be re-sown if it is well managed. For best results, do not graze first year stands during flowering. After seed set, remove plant residues in late summer to promote better seedling regeneration. *Zulu II* suits is tolerant of moderately acidic soils. Sow treated seed.

Suggested seed mixes

Dairy		kg/ha
12 month high performance crop	<i>Tabu</i> + Italian ryegrass	18-22
	<i>Laser</i> Persian clover	4
	<i>Vista</i> balansa clover	3
	<i>Morrow</i> red clover (coated)	6
	Total	31-35
Chicory/annual clover crop	<i>501 Chicory</i>	8
	<i>Zulu II</i> arrowleaf clover	8
	Total	16
Sheep, Beef and Deer		
8-10 month pure clover sward	<i>Zulu II</i> arrowleaf clover	10
Hill country oversow mix	<i>Safin</i> cocksfoot	8
	<i>Weka</i> white clover	2
	<i>Apex</i> white clover	2
	Sub clover	6
	<i>Zulu II</i> arrowleaf clover	4
Total	22	



Zulu II Arrowleaf Clover is owned and marketed by Barenbrug Agriseeds
Zulu II Arrowleaf Clover is protected under the NZ Plant Variety Rights Act 1987

VISTA

BALANSA CLOVER

Vista is mid-late flowering and produces high quality feed in winter and early spring. It has excellent tolerance to waterlogging and is an ideal short term crop option for grazing, silage or hay particularly on wet or poorly drained soils.

Fills the gap

Vista was selected for improved growth during winter and early spring to help fill the typical early feed deficit on many farms.

System fit

Vista will significantly improve dairy farm feed quality and early season production when autumn-sown with annual ryegrass for winter grazing. This mix will also make high quality silage. A straight sward of *Vista* is excellent feed for lactating ewes and growing lambs before the paddock goes into spring-sown crop.

Management

Vista is very adaptable, and can handle a range of different grazing systems. For maximum animal performance and pasture regrowth, graze down to 4-5 cm residual.

Suggested seed mixes

Dairy		kg/ha
Winter ryegrass crop	<i>Hogan</i> annual ryegrass	25-30
	<i>Vista</i> balansa clover	3
	<i>Laser</i> Persian clover	4
	Total	32-37
Sheep, Beef and Deer		
7-9 month pure finishing sward	<i>Vista</i> balansa clover	6
Winter oat crop	<i>Hattrick</i> oats	80
	<i>Vista</i> balansa clover	4
	Total	84



Vista balansa clover is ideal for grazing, silage or hay.

LASER

PERSIAN CLOVER

Late-flowering *Laser* produces valuable high quality feed from winter through early summer, for improved animal production and finishing. It is soft seeded, very fast establishing, and moderately tolerant of water logging.

Later growth

Laser has a different growth curve to *Vista* and *Zulu II*, flowering 30 days later, so extending growth into summer.

System fit

On dairy farms, add *Laser* to short-term pastures to improve feed quality and extend DM and animal production from early spring to early summer. For sheep and beef, it is an excellent option to increase liveweight gains. *Laser* also suits hay/silage making.

Management

Laser can be grazed down to residuals of 2-3 cm during winter. In spring, rotationally graze to residuals of 4-5 cm to maximise animal performance, and pasture regrowth. Avoid over grazing, which will remove developing stems.

Conditions

Laser can tolerate mild salinity, cold temperatures and partially waterlogged soils. It is susceptible to slugs and springtails during establishment. Use treated seed, slug bait if needed, and include an insecticide at spray out. *Laser* is resistant to clover scorch.

Suggested seed mixes

Dairy		kg/ha
12-18 month high performance crop	<i>Tabu</i> + Italian ryegrass	18-22
	<i>Laser</i> Persian clover	4
	<i>Vista</i> balansa clover	3
	<i>Morrow</i> red clover (coated)	6
	Total	31-35
6-8 month winter crop	<i>Hogan</i> annual ryegrass	25-30
	<i>Laser</i> Persian clover	4
	<i>Vista</i> balansa clover	3
	Total	32-37
Sheep, Beef and Deer		
8-10 month pure finishing sward	<i>Laser</i> Persian clover	10
Two year finishing crop	<i>Captain</i> plantain	10
	<i>Laser</i> Persian clover	4
	<i>Vista</i> balansa clover	3
	<i>Morrow</i> red clover (coated)	6
	<i>Weka</i> white clover	4
Total	27	



Late-flowering Laser Persian clover improves animal production and finishing.

501

CHICORY

501 Chicory is best suited as a 6-8 month summer crop, with rapid establishment and excellent re-growth, to give very high DM yield and an extra grazing over some other cultivars. It also has an erect growth habit, which provides high utilisation.

Avoid FEI milk penalties

The FEI (Fat Evaluation Index) milk grading system came into effect from in 2018. Industry trials have shown chicory has no effect on the FEI, whereas DairyNZ's rule of thumb is to feed less than 3kg DM/cow/day of PKE to avoid milk penalties.


Rapid establishment

501 Chicory establishes very quickly, meaning less down time before the first grazing. In trials and on-farm across a wide range of different soil types *501 Chicory* has stood out right from the start, particularly in challenging dry conditions.

Excellent DM yield

501 Chicory's extra yield is predicted to produce an extra 7% kg MS, giving an additional income of \$325/ha (based on \$6.50/kg MS) over *Choice* chicory.

Modelled Chicory MS production (Relative to trial mean = 100).



Data based on yield info from the combined trial analysis of Cambridge 11-12, and Canterbury 12-13. 2 trial lines have been removed from the graph. Assumptions used were: ME of chicory is 12 MJ ME/kg DM and 132 MJ ME to produce 1kgMS.

Advantage of 501 + Morrow

The combination of *501 + Morrow* red clover or *501 + Zulu II* arrowleaf clover performs well. Like *501*, these clovers are deep rooted giving them a significant advantage in summer dry conditions. These clovers grow well between the chicory plants filling gaps often otherwise taken up by weeds. They also fix nitrogen reducing fertiliser requirements for the crop.

High ME

Both chicory and red clover are highly palatable to livestock and are both high in ME. During summer dry conditions, *501 Chicory* and *Morrow* red clover will maintain an ME of around 12, whereas ryegrass pastures generally maintain an ME of 8.5-10.5.

Environmental gains

501 Chicory offers a range of important environmental benefits: It doesn't need insecticide sprays (unlike brassica crops); Its up to 1.5m deep tap root improves soil structure; Mines deep soil N and can recover excess soil N left after winter-grazed crops; Research has shown heifers grazing chicory urinated more frequently without increasing urinary output, or urinary N, potentially reducing N loading and subsequent nitrate leaching from soil, and; Facial eczema spores are much lower than on ryegrass pastures, reducing the need to intervene with your animals.

Management

Sow chicory into a firm, fine, weed-free seed bed where soil temperatures are consistently above 12°C in spring. Roll before and after sowing to help get a uniform germination. First grazing should occur when plants reach the seven leaf stage. Targets for grazing are:

- Pre-grazing targets: 3000 kg DM/ha or 25-35 cm height.
- Post-grazing residual target: 5 cm.

How many ha?

For dairy farms, sow 3 ha of *501 Chicory* per 100 cows to provide 3 kg DM of chicory/cow/day. Area to be sown can be calculated from the following table:

Chicory/cow to be fed	Area of chicory to sow	Daily area of chicory*
2 kg DM/day	2 ha/100 cows	0.1 ha/100 cows
3 kg DM/day	3 ha/100 cows	0.15 ha/100 cows
4 kg DM/day	4 ha/100 cows	0.2 ha/100 cows

*Assuming 21 day grazing rotation.

When to resow pasture

501 Chicory will look great going into autumn. In spite of this it is more important to get new pasture established early, rather than continue to graze chicory into late autumn.

Suggested seed mixes

Use		kg/ha
For a chicory crop	<i>501 Chicory</i>	8-10
	Total	8-10
	Chicory/red clover crop	
	<i>501 Chicory</i>	6-8
	<i>Morrow red clover</i>	4
	Total	10-12
Chicory/annual clover crop	<i>501 Chicory</i>	8
	<i>Zulu II arrowleaf clover</i>	8
	Total	16



501 Chicory is owned and marketed by Barenbrug Agriseeds
501 Chicory is protected under the NZ Plant Variety Rights Act 1987

CAPTAIN CSP


PLANTAIN

We've called *Captain* a 'cool season plantain (CSP)' due to its extra growth in this period, the most valuable feed in farm systems, with environmental advantages too. *Captain* also has excellent summer yield and drought tolerance.

Outstanding cool season production

Outstanding cool season production: *Captain CSP* yields significantly more through the cool season period as shown in the graph below. Plantains vary hugely in winter growth, as you can see in the photo.

Cool season DM yield data combined from three one year dryland Canterbury trials sown between 2013 – 2018*



* Cool season yield is a total of autumn, winter & early spring periods. LSD (5%) lettering given on yield bars, cultivars with the same letter are not significantly different. ¹ Provisional rating, cultivar has only been in one trial.

Reduced N leaching

Initial investigations indicate that plantain can mitigate N leaching via a number of mechanisms, including direct activities on soil N mineralisation and direct uptake of N through growth. The greater cool season activity of *Captain CSP* will enhance both of these mechanisms when it is most needed, as N leaching mainly happens when soils are wet through the late autumn, winter and early spring.



Captain CSP (centre) showing its significant cool season yield advantage over other cultivars at the end of July at Courtenay 190m a.s.l.

High total DM yield

Captain CSP also produces strongly across the other seasons too. It's deep rooting, with high summer yield providing additional protein and feed quality over the warmer months, particularly in summer dry areas.

Animal performance

Plantain is easily digestible, improving stock appetite especially over dry summer months when grasses are of lower feed quality. It is also higher in essential minerals like P, K, S, Ca, Mg, Na, Zn, Cu, B and Co than ryegrass/clover pastures.

Plant type

Captain CSP is a distinctive narrow-leaved plant with upright growth habit for high utilisation. It has a deep, fibrous root system, and good compatibility with other species. It has good persistence, and can last three years under good management.

Sheep, beef, deer systems

Captain CSP can be used as a high LWG finishing crop, for example mixed with red, white, and annual clovers. Here the annual clovers (Persian, arrowleaf) provide most of the LWG through the first year, with red and white clovers providing it after that.

Dairy systems

Captain CSP can be used as a summer crop, or sown as part of a pasture mix at 2-4 kg/ha to increase summer feed quality in dryland situations. *Captain* can also be used as part of a specialist high-yielding, quality 2-3 year pasture, with *Shogun NEA* hybrid ryegrass and *Kotuku* white clover.

Suggested seed mixes

Sheep, Beef, Deer		kg/ha
Two year high LWG finishing crop	<i>Captain</i> plantain	10
	<i>Laser</i> Persian clover	4
	<i>Vista</i> balansa clover	3
	<i>Morrow</i> red clover (coated)	6
	<i>Weka</i> white clover	4
Total		27
Perennial pasture mix	<i>Tyson</i> or <i>Rohan SPR</i> ryegrass	18
	<i>Safin</i> cocksfoot	4
	<i>Weka</i> white clover	4
	<i>Morrow</i> red clover	4
	<i>Captain</i> plantain	2
Total		32
Dairy		kg/ha
Perennial pasture mix	<i>Trojan</i> or <i>Governor</i> ryegrass	22
	<i>Kotuku</i> white clover	2
	<i>Weka</i> white clover	2
	<i>Captain</i> plantain	2
Total		28
Specialist 2-3 year pasture	<i>Shogun NEA</i> hybrid ryegrass	30
	<i>Kotuku</i> white clover	4
	<i>Captain</i> plantain	2
Total		36

INVITATION

SWEDE

Invitation is a late maturing, yellow-fleshed swede, with very high bulb and leaf yield. It provides winter feed with excellent animal health for sheep, cattle and deer.

High yield & disease tolerance

Invitation produces excellent total DM yields with good dry rot tolerance and resistance to club root and powdery mildew. *Invitation* is not recommended as a second crop.

Total DM yield, dry rot tolerance and club root infection level.

Cultivar	Total DM yield*		Dry rot tolerance**		Club root***	
	(Trial mean =100)		% of bulbs not infected	% bulbs badly infected	% of bulbs not infected	
<i>Invitation</i>	112	a	57	a	5	a
<i>Aparima Gold</i>	103	b	36	ab	11	a
<i>Major Plus</i>	96	c	10	bc	56	b
<i>Dominion</i>	92	c	6	c	71	b
<i>Domain</i> ◊	74	d	NT	NT	NT	NT
Trial mean	12.6 t DM/ha		21%		41%	
					60%	

*From 8 Southland trials, from 2006/07 to 2011/13. **From a Southland trial in 2008/09 under moderate to high dry rot pressure in a 2nd crop paddock. *** From a Southland trial in 2010/11 under moderate to high club root pressure in a 2nd crop paddock. NT = Not tested. Statistical significance lettering given for 5% LSD level, cultivars with same letter are not significantly different. ◊ = Provisional results. *Domain* was in 2 of the 8 trials.

Late flowering

Invitation is very late flowering, meaning the crop stays vegetative longer into spring than other cultivars. This minimises the chance of animal health problems associated with 'bolting' swede crops, as seen in spring 2014 in Southland. No issues were reported on *Invitation* swedes.

Swede flowering scores*

Cultivar	Flowering score
<i>Invitation</i>	7.2 a
<i>Major Plus</i>	6.7 ab
<i>Domain</i>	6.5 ab
<i>Dominion</i>	4.8 c
<i>HT Swede</i>	3.4 d
<i>Aparima Gold</i>	3.1 d
Trial mean	6.1

*Results from 2 trials in Southland sown 2008 and 2012. Statistical significance lettering given for 5% LSD level, cultivars with the same letter are not significantly different. Scored on a 1 - 9 basis. Where 1 = full flowering swede crop, 4 = stem elongation, green seed head appeared, 7, small degree of elongation, 9 = no sign of stem elongation.

Good leaf yield

Invitation produces high leaf yields showing a significantly higher leaf percentage than other cultivars in trials. This lifts the overall protein level of the crop and is helpful when introducing swedes into an animal's diet, particularly for younger stock.

Bulb & leaf keeping

Invitation has shown high bulb keeping ability and leaf retention in trials, helping maintain its feed quality and quantity through to the end of winter.

Using Invitation

Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	
Sow							Graze			
Maturity date:			170-250 days							
Typical yield:			10-18 t DM/ha (depending on season)							
ME:			12-14 MJ/kg DM							
Sowing rate:			0.5-0.8 kg/ha ridged 0.8-1.5 kg/ha drilled							



Invitation produces a higher proportion of leaf than other varieties, providing more protein.

INTERVAL

RAPE

Interval is a tall, fast establishing rape ideal for summer, autumn and winter feed. It offers very high yield for all stock types.

Interval is a rape-kale cross, giving high yield with regrowth ability.

Flexible sowing date

Interval can be sown from spring through to early autumn to provide a bulk of high quality feed typically in 90-110 days. Spring sowings can be grazed in summer/early autumn then left to regrow for winter feed.

High yield

Interval has performed well in trials, providing excellent DM yield.

Total winter DM yield*

Cultivar	Trial mean = 100%	
<i>Interval</i>	126	a
<i>Goliath</i>	125	a
<i>Greenland</i>	118	a
<i>Winfred</i>	92	b
<i>Titan</i>	88	b
Trial mean (t DM/ha)	5.3	

*Results from 2 trials in Canterbury during 2008 and 2009 (February sown, June/July harvested). Statistical significance lettering given for 5% LSD level, cultivars with the same letter are not significantly different.

Winter utilisation

Compared to kale, rape typically has higher stem feed quality, and is better utilised by stock.

Other characteristics

Interval has excellent tolerance of dry conditions. It also has strong frost tolerance and resistance to powdery mildew.

Using *Interval*

	Sept	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	
		Sow											
						Graze							
Maturity date:	90-110 days												
Typical yield:	5-8 t DM/ha (depends on sowing time & no. of grazings)												
Typical ME:	12 MJ/kg DM												
Sowing rate:	4 kg/ha												

DYNAMO

TURNIP

Dynamo turnip is a high yielding summer crop which is ideal for dairy cows. It provides large volumes of low cost quality feed to help maintain milk production when pasture quality and quantity declines.

DM yield

In trials *Dynamo* has shown high yield, not significantly different from the other top cultivars.

Total DM yield*

Cultivar	Trial mean = 100%
<i>Barkant</i>	110 a
<i>Dynamo</i>	103 ab
<i>Marco</i>	102 ab
<i>Rival</i>	102 ab
<i>Envy</i> [◇]	101 ac
<i>White Star</i>	98 bc
<i>Green Globe</i>	93 c
Trial mean (t DM/ha)	8.2

* From 10 trials in Waikato (7), Taranaki (2) & Canterbury (1) from 2006/07 to 2008/09. ◇ = Provisional result: *Envy* was only in 2 of the 10 trials. Statistical significance lettering given for 5% LSD level, cultivars with the same letter are not significantly different.

Low cost summer feed

Sowing a poor performing pasture in *Dynamo* makes financial sense. It can provide feed for around 20 c/kg DM*.

*Turnips for 20 c/kg DM - assumptions:

- Turnip crop yield 11.5 t DM/ha, with 12 ME.
- 5.5 t of old pasture growth is forgone while the paddock is in crop.
- Cost of growing crop = \$1,200/ha (spray out plus insecticide, full cultivation, fertiliser, treated seed, slug bait, two post emergence herbicides/insecticides).
- \$1,200/6,000 kg DM extra yield = 20 c/kg DM

High bulb percentage

Summer turnips produce their yield in different ways. *Dynamo* produces a good level of bulb (around 48% of total yield), giving it an advantage in seasons when high levels of leaf diseases or pests are present.

Using *Dynamo*

	Oct	Nov	Dec	Jan	Feb	Mar
Sow						
Graze						
Maturity date:	60-90 days					
Typical yield:	8-16 t DM/ha (depending on season)					
ME:	12 MJ/kg DM					
Sowing rate:	2-3 kg/ha					

FODDER BEET

Fodder beet types

It's important to choose the correct fodder beet variety for your feed requirements and intended use (grazing, lifting or both). Good starting points for this decision are bulb DM content, and whether the crop is only intended to be lifted. Fodder beet can be largely divided into three groups based on these factors:

Low bulb DM% (12-15%)

Lower yield potential, usually with a high % of bulb above ground (50%+). Only suited to grazing in situ.

Medium-high bulb DM% (16-20%)

Higher yield potential than low DM % types, and can be grazed in situ e.g. *Robbos*. Some can also be successfully lifted or grazed

Lifting types

Bulbs sit lower in the ground, generally not suitable for grazing in situ. Very high DM % types (e.g. *Blizzard*) are best for maximum yield potential and increased storage life.

System fit

Thanks to its ability to grow a large volume of high quality, high utilisation feed that can be used from autumn to spring, fodder beet suits several different farm systems. Its high yield potential also frees up land for other uses, which is a major plus. Alternatively you can increase daily allowances for improved live weight gains.

This crop provides flexible winter grazing and can also be used to extend dairy cow lactation by either grazing in situ or lifting and feeding to stock on pasture. Successful grazing entails correct stock transition.

For further information see Barenbrug Agriseeds Fodder Beet Product Guide or download the guide from www.agriseeds.co.nz

	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	July	Aug	Sep
Robbos												
Dairy	Precision sown.						Extend lactation, start winter transition.		Winter feed.		Supplement spring pasture covers.	
Beef/Sheep/Deer	Precision sown.						High ME feed for liveweight gain or maintenance from autumn to spring.					
Blizzard												
Lifting fodder beet	Precision sown.						Mechanically lifted and fed to stock for a high ME supplement from autumn through to early summer.					
Maturity:	Once herbicide withholdings are met. 170 days+ to maximise yield.											
Typical Yield	18-24 t DM/ha average. 25 t DM/ha+ possible with good summer moisture and fertility.											
Sowing rate:	80,000 seeds/ha grazing. 100,000 seeds/ha lifting.											

ROBBOS


FODDER BEET

High DM yield

Robbos has been an excellent, consistent performer with high DM yield and improved leaf holding ability to provide more protein and better animal nutrition.

The medium DM content of *Robbos* makes it capable of producing a 4.4t DM/ha higher DM yield than lower DM types such as *Brigadier* worth an extra \$1320/ha (assuming a 30c/kg DM value for high ME winter feed). Its palatable orange-yellow bulbs are suitable for grazing by all stock types.

Fodder beet DM yields - low and medium DM cultivars



*Combined analysis of 5 trials from 2014-2017, varieties in two or more trials are presented. Cultivars with the same statistical significance letter are not significantly different at the LSD 5% level.

Higher protein = less silage



Robbos (middle 3 rows) showing excellent leaf holding ability versus *Kyros* (left) and *Enermax* (right) in Canterbury trial.

In 2018 trials *Robbos* leaf tested at 24.5% protein, significantly higher than *Feldherr*, *Brigadier*, *Monro* and *SF1505* (which averaged 21%). This increased protein from *Robbos* is equivalent to 4.5t/ha of good pasture silage*, which could save \$350/ha based on \$0.40/kg DM for good pasture silage.

*23% DM, 17% protein - based on Feed Composition for Good Pasture Silage, DairyNZ Facts and Figures


BLIZZARD FODDER BEET

Blizzard is a white-skinned fodder beet with excellent leaf holding ability and disease resistance, which helps to maximise yield potential. It should be used when maximum yield/ha is sought from a lifted crop, and has 20-22% DM content.

High DM yield

Blizzard has performed very well in trials, with significantly higher total DM yield than low - medium DM cultivars. *Blizzard* is not recommended for grazing because it has a high proportion of the bulb in the ground, reducing utilisation.

Fodder beet combined DM yield analysis. Data from 7 trials over 2008-2018*.



* Combined data of seven trials from 2008-2018: Canterbury (4), Southland (3). Statistical significance lettering (LSD 5%) given. Cultivars with the same letter are not significantly different.

Storage

The higher DM content of *Blizzard* enables it to be kept for longer in a windrow after being lifted. When leaves are removed. *Blizzard* can be stored for up to 5 months, but bulbs will dry out over time, so DM % may need to be re-tested to allow for accurate feed allocation.

Using *Blizzard*

Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep
Precision sown									Mechanically lifted and fed to stock for a high ME supplement from autumn through to spring		
Feeding method:			Lifting only								
Typical yield:			20-25 t DM/ha average; > 26 t DM/ha with summer moisture								
Typical ME:			12-13 MJ/ME								
Sowing rate:			100,000 seeds/ha								

INSECT CONTROL RATING FOR ENDOPHYTES

Summary

These ratings are indicative and may vary slightly between cultivars. If Argentine stem weevil or black beetle are present at sowing, an appropriate seed treatment is recommended to improve insect resistance during establishment. The ratings in this table are based in part on glasshouse studies where test plants are 100% infected with endophyte, whereas commercial seed must meet minimum standards of 70% of seeds infected. These tables were compiled by AgResearch, Agricom, Barenbrug Agriseeds, Cropmark, Grasslanz and PGG Wrightson Seeds.

Endophyte insect control for perennial ryegrass, festulolium & short-term (hybrid) ryegrass.

	Argentine stem weevil	Pasture mealy bug	Black beetle adult	Root aphid	Porina	Grass grub	Field cricket
Diploid perennial ryegrass							
AR1	◆◆◆◆	◆◆◆◆	◆	- ²	-	-	Not tested
NEA2	◆◆◆	(◆◆◆◆)	◆◆◆	◆◆	Not tested	-	Not tested
AR37	◆◆◆◆ ¹	◆◆◆◆	◆◆◆	◆◆◆◆	◆◆◆	◆	Not tested
SE	◆◆◆◆	◆◆◆◆	◆◆◆	◆◆	◆	-	Not tested
WE	-	-	-	-	-	-	Not tested
Tetraploid perennial ryegrass							
AR1	(◆◆◆)	(◆◆◆◆)	◆	- ²	-	-	Not tested
AR37	(◆◆◆) ¹	(◆◆◆◆)	◆◆◆	◆◆◆◆	(◆◆◆)	◆	Not tested
WE	-	-	-	-	-	-	Not tested
Festulolium							
U2	◆◆◆◆	(◆◆◆◆)	◆◆◆◆ ³	◆◆◆◆	(◆◆)	◆◆◆	◆◆◆
Italian and short term (hybrid) ryegrass							
AR1	◆◆	(◆◆◆◆)	◆	- ²	Not tested	-	Not tested
NEA	Not tested	(◆◆◆◆)	◆◆◆	Not tested	Not tested	-	Not tested
AR37	◆◆◆ ¹	(◆◆◆◆)	◆◆◆	Not tested	Not tested	-	Not tested
WE	-	-	-	-	-	-	Not tested

Notes on table:

¹ AR37 endophyte controls Argentine stem weevil larvae, but not adults. While larvae cause most damage to pastures, adults can damage emerging grass seedlings. In Argentine stem weevil prone areas it is recommended to use treated seed for all cultivars with novel endophyte.

² AR1 plants are more susceptible to root aphid than plants *Without* endophyte.


³ Also active against black beetle larvae

Key to tables

-	No control.
◆	Low level control: Endophyte may provide a measurable effect, but is unlikely to give any practical control.
◆◆	Moderate control: Endophyte may provide some practical protection, with low to moderate reduction in insect population.
◆◆◆	Good control: Endophyte markedly reduces insect damage under low to moderate insect pressure. Damage may still occur when insect pressure is high.
◆◆◆◆	Very good control: Endophyte consistently reduces insect populations and keeps pasture damage to low levels, even under high insect pressure.
()	Provisional result: Further results needed to support the rating. Testing is ongoing.

Endophytes control Argentine stem weevil differently

NEA endophytes and *ARI* control adult ASW, preventing their feeding and egg laying, so protecting pastures. *AR37* gives no control of adult ASW, but controls ASW larvae, stopping larval feeding, so protecting pastures.



ENDOPHYTE ANIMAL SAFETY

Summary

These ratings are indicative. Animal performance and health can vary under different management systems and between seasons.

Sheep & lambs

	<i>AR1</i>	<i>NEA</i>	<i>NEA2</i>	<i>AR37</i>	<i>U2</i>	<i>Standard endophyte</i>	<i>Without endophyte</i>
Freedom from ryegrass staggers	◆◆◆◆	◆◆◆◆	◆◆◆◆	◆◆◆◆ ²	◆◆◆◆	◆ ¹	◆◆◆◆
Animal production	◆◆◆◆	◆◆◆◆	◆◆◆◆	◆◆◆◆ ³	◆◆◆◆	◆◆ ¹	◆◆◆◆

Notes on sheep & lambs table:

¹*Standard* endophyte can cause severe ryegrass staggers, can significantly decrease lamb growth rates in summer and autumn, and significantly increase dags.

²Ryegrass containing *AR37* endophyte can cause severe ryegrass staggers, but the frequency of ryegrass staggers is much lower than for ryegrass with *Standard* endophyte. *One50 AR37* may give rise to higher instances of ryegrass staggers than other *AR37* cultivars in some situations.

³Lambs grazing ryegrass containing *AR37* endophyte can have reduced LWG during periods of severe staggers.

Dairy cows & beef cattle

	<i>AR1</i>	<i>NEA</i>	<i>NEA2</i>	<i>AR37</i>	<i>U2</i>	<i>Standard endophyte</i>	<i>Without endophyte</i>
Freedom from ryegrass staggers	◆◆◆◆	◆◆◆◆	◆◆◆◆	◆◆◆◆ ²	◆◆◆◆	◆◆ ¹	◆◆◆◆
Animal production	◆◆◆◆	Not tested	Not tested	◆◆◆◆ ³	◆◆◆◆	◆◆◆ ¹	◆◆◆◆

Notes on dairy cows & cattle table:

¹*Standard* endophyte can cause ryegrass staggers, and has been shown to depress milksolids production through summer and autumn.

²While ryegrass staggers has not been observed on cattle and dairy cows, it could occur on rare occasions.

³In dairy trials overall MS production from ryegrass containing *AR37* endophyte is not significantly different from that with *AR1*. A small reduction in MS was observed over summer on ryegrass containing *AR37*. A contributing factor to this was the lower clover content in *AR37* pastures.

Key to tables

-
- ◆◆ **Moderate animal performance and health:** This endophyte is known to regularly cause significant problems.

 - ◆◆◆ **Good animal performance and health:** This endophyte can cause problems from time to time.

 - ◆◆◆◆ **Very good animal performance and health.**



Member of the Royal Barenbrug Group

Barenbrug® Agriseeds® are registered trademarks of New Zealand Agriseeds Limited.

* *Trojan* is a perennial ryegrass certified under the New Zealand seed certification scheme as *lolium boucheanum*.

0800 449 955

www.agriseeds.co.nz

**COMMITTED TO INNOVATION.
COMMITTED TO FARMING.**