





Storm, as with many white clovers, is susceptible to a wide range of diseases that include Alfalfa Mosaic Virus (*Alfamovirus* spp.), white clover Mosaic Virus (*Potexvirus* spp.) and several Root Rots caused by various fungal pathogens.

Variety Management/Agronomy:

White clover is generally sown as a component of a pasture mix with grasses. For sowing a pure stand multiply sowing rates by 2 - 3 times. White clover is tolerant of, and will persist under a wide range of management systems and has high feed value. Its ability to fix atmospheric nitrogen makes a substantial contribution to the growth of companion grasses. White clover will grow over a wide range of soil and fertility conditions although a pH of 5.4 or higher with reasonable phosphorus levels is required for optimal results. It has poor tolerance of drought conditions and is best suited to medium-high rainfall or irrigation, where it will respond well to spring and summer moisture.





Key Features:

- Australian bred white clover
- Mid-maturity
- · High production in winter and summer
- Tall plant type that can aggressively compete in a mixed sward with ryegrass
- Excellent seedling vigour and is quick to establish
- Excellent all season growth with very high yield potential across all seasons
- Stolen density high compared to other large leaf types
- Persistent under cutting and remains dense

Description:

Storm white clover is a tall plant type that can aggressively compete and actively grow up through the sward. It offers excellent total forage production with exceptional yield potential across all seasons. Fast to establish, Storm is quick out of the ground, providing better competition with grasses and the potential to spray broadleaf weeds earlier. It also is persistent under grazing with high stolon density for a large leaf type clover. High production in winter and summer.

Product Fit:

High performance, high output systems in the irrigation and higher rainfall regions – suited to use where other white clovers are used. May not be suited to very tight grazing under sheep.

Breeding History:

Storm is an Irrigation by Tamar cross bred by the Victorian DPI. The key selection criteria included persistence, high production, with a specific focus on winter growth, leaf size and high stolon density.


Pest Resistance:


Storm is susceptible to Red Legged Earth Mites (*Halotydens destructor*) and control measures will be required either prior to, or soon after germination. Other insects that need to be monitored carefully include Pea Aphid (*Acyrthosiphon pisum*), Blue Oat Mites (*Penthaleus major*) and Cut Worms (*Agrotis munda*).




Trial Results:

Storm has excellent total forage yield with exceptional yield potential across all seasons. Storm was the highest yielding variety in the trial for total yield. Seasonally it was the highest yielding variety in all seasons other than winter where it was only fractionally behind Haifa.





Trial data: Heritage Seeds Howlong NSW. Sown spring 2009.



^{*} Base data for graph produced by AgResearch. Storm and Haifa's position estimated using stolon growing point and leaf size measurements taken at Howlong by Heritage Seeds in 2013. Weka's position estimated using growing point density and leaf size measurements by Agriseeds 2005-08.

Plant Breeders Rights (PBR): This variety is registered under Plant Breeders Rights (PBR) in Australia. Unauthorised commercial propagation or any sale, conditioning, export, import or stocking of propagating material is an infringement under the Plant Breeders Rights Act (1994). Any breach of this legislation will leave the grower liable for prosecution.

Disclaimer: The information presented in this brochure is from official and other sources and is considered to be reliable. It is provided in good faith and every care has been taken to ensure its accuracy. Heritage Seeds does not accept any responsibility for the consequences that may arise from the acceptance of recommendations or the suggestions made.

